

# A Theorist's View on the Current Status of SUSY Searches

David Shih  
Rutgers University

BNL Workshop on SUSY with 5/fb

May 3, 2012

# SUSY at the LHC

# SUSY at the LHC

- The LHC has searched for SUSY far and wide. Nothing but limits so far.
- What are the implications for the underlying model of SUSY-breaking and mediation?
- .....

# SUSY at the LHC

- The LHC has searched for SUSY far and wide. Nothing but limits so far.
- What are the implications for the underlying model of SUSY-breaking and mediation?
- .....
- Has SUSY been excluded?

# SUSY at the LHC

- The LHC has searched for SUSY far and wide. Nothing but limits so far.
- What are the implications for the underlying model of SUSY-breaking and mediation?
- .....
- Has SUSY been excluded?
- Is it time to throw in the towel?

# SUSY at the LHC

- The LHC has searched for SUSY far and wide. Nothing but limits so far.
- What are the implications for the underlying model of SUSY-breaking and mediation?
- .....
- Has SUSY been excluded?
- Is it time to throw in the towel?
- Why are we here?

# SUSY at the LHC

- The LHC has searched for SUSY far and wide. Nothing but limits so far.
- What are the implications for the underlying model of SUSY-breaking and mediation?
- .....
- Has SUSY been excluded?
- Is it time to throw in the towel?
- Why are we here?
- Should we all go home?

# SUSY at the LHC



# SUSY at the LHC

- In this talk, I will review the current status of SUSY searches and their implications for the underlying model.
- My remarks are meant to represent a purely personal point of view.
- ....

# SUSY at the LHC

- In this talk, I will review the current status of SUSY searches and their implications for the underlying model.
- My remarks are meant to represent a purely personal point of view.
- ....
- While it is disappointing that SUSY was not “just around the corner,” I hope you will agree that there is still much discovery potential remaining!

# The SUSY Paradigm



MSSM

$M_{weak}$

SUSY cannot be broken  
directly in the MSSM.

# The SUSY Paradigm

SUSY Sector

$$\sqrt{F}$$


Rather, it must first be broken  
in a “hidden sector.”

MSSM

$$M_{weak}$$


SUSY cannot be broken  
directly in the MSSM.

# The SUSY Paradigm




# The SUSY Paradigm

There are many possibilities for a complete SUSY model




# The SUSY Paradigm

There are many possibilities for a complete SUSY model



# The SUSY Paradigm


There are many possibilities for a complete SUSY model






# The SUSY Paradigm

There are many possibilities for a complete SUSY model




# The SUSY Paradigm

There are many possibilities for a complete SUSY model




# The SUSY Paradigm

There are many possibilities for a complete SUSY model




# The SUSY Paradigm

There are many possibilities for a complete SUSY model




# The SUSY Paradigm

There are many possibilities for a complete SUSY model



# The SUSY Paradigm


There are many possibilities for a complete SUSY model






# The SUSY Paradigm

There are many possibilities for a complete SUSY model



# The SUSY Paradigm


There are many possibilities for a complete SUSY model






# The SUSY Paradigm

There are many possibilities for a complete SUSY model




# The SUSY Paradigm

There are many possibilities for a complete SUSY model




# The SUSY Paradigm

There are many possibilities for a complete SUSY model



# The SUSY Paradigm

There are many possibilities for a complete SUSY model



# SUSY Scenarios

Type	Mediation Scale	LSP	Pros	Cons
Gravity mediation	Planck-scale	Neutralino or sneutrino	WIMP DM candidate; automatic $\mu/B\mu$	severe SUSY flavor problem; uncalculable framework
Anomaly mediation	super-Planck-scale	Neutralino (wino)	no SUSY flavor problem	tachyonic sleptons; requires “sequestering”
Gauge mediation	sub-Planck-scale	gravitino	no SUSY flavor problem; calculable framework; viable spectrum	no WIMP DM $\mu/B\mu$ problem

# Two views of the SUSY-breaking Scale



# Two views of the SUSY-breaking Scale



The scale of SUSY breaking determines the mediation mechanism.



# Two views of the SUSY-breaking Scale



The scale of SUSY breaking determines the mediation mechanism.

Gauge  
mediation






# Two views of the SUSY-breaking Scale




The scale of SUSY breaking determines the mediation mechanism.



# Two views of the SUSY-breaking Scale




The scale of SUSY breaking determines the mediation mechanism.



# Two views of the SUSY-breaking Scale



The scale of SUSY breaking determines the mediation mechanism.




It also determines the behavior of the lightest MSSM superpartner (LOSP).

# Two views of the SUSY-breaking Scale



The scale of SUSY breaking determines the mediation mechanism.




It also determines the behavior of the lightest MSSM superpartner (LOSP).

# Two views of the SUSY-breaking Scale



The scale of SUSY breaking determines the mediation mechanism.




It also determines the behavior of the lightest MSSM superpartner (LOSP).

# Two views of the SUSY-breaking Scale



The scale of SUSY breaking determines the mediation mechanism.




It also determines the behavior of the lightest MSSM superpartner (LOSP).

# Two views of the SUSY-breaking Scale



The scale of SUSY breaking determines the mediation mechanism.




It also determines the behavior of the lightest MSSM superpartner (LOSP).

# Two views of the SUSY-breaking Scale




The scale of SUSY breaking determines the mediation mechanism.



It also determines the behavior of the lightest MSSM superpartner (LSP).


Viewed like this, there is no material difference between high-scale GMSB, gravity mediation, and anomaly mediation!!!





Inclusive SUSY collider signatures are mainly dictated by:

- Production mechanism (strong or EW superpartner)
- Identity of the LOSP
- Lifetime of the LOSP





Inclusive SUSY collider signatures are mainly dictated by:



- Production mechanism (strong or EW superpartner)
- Identity of the LOSP
- Lifetime of the LOSP

We will now survey some of the possibilities, keeping in mind the underlying mediation mechanism.

# SUSY production at the LHC





# SUSY production at the LHC





- For EW production, LHC7 @ 1/fb  $\sim$  Tevatron @ 10/fb

# SUSY production at the LHC





- For EW production, LHC7 @ 1/fb  $\sim$  Tevatron @ 10/fb
- For strong production, LHC7 @ 1/fb  $\gg$  Tevatron @ 10/fb

# SUSY production at the LHC





# SUSY production at the LHC



“Kinematic reach” of LHC7:



# SUSY production at the LHC



“Kinematic reach” of LHC7:  $M_{\text{wino}} \sim 700$  GeV





# SUSY production at the LHC



“Kinematic reach” of LHC7:  $M_{\text{wino}} \sim 700$  GeV

$M_{\text{gluino}} \sim 1200$  GeV

# SUSY production at the LHC





“Kinematic reach” of LHC7:  $M_{\text{wino}} \sim 700$  GeV

$M_{\text{gluino}} \sim 1200$  GeV

“Kinematic reach” of LHC14:

# SUSY production at the LHC




“Kinematic reach” of LHC7:  $M_{\text{wino}} \sim 700$  GeV

$M_{\text{gluino}} \sim 1200$  GeV


“Kinematic reach” of LHC14:  $M_{\text{wino}} \sim 1600$  GeV

# SUSY production at the LHC



“Kinematic reach” of LHC7:  $M_{\text{wino}} \sim 700$  GeV


“Kinematic reach” of LHC14:  $M_{\text{wino}} \sim 1600$  GeV



$M_{\text{gluino}} \sim 1200$  GeV


$M_{\text{gluino}} \sim 2500$  GeV

# SUSY production at the LHC



“Kinematic reach” of LHC7:  $M_{\text{wino}} \sim 700$  GeV

“Kinematic reach” of LHC14:  $M_{\text{wino}} \sim 1600$  GeV




$M_{\text{gluino}} \sim 1200$  GeV

$M_{\text{gluino}} \sim 2500$  GeV

Yardsticks to measure the current progress

# CMS Preliminary

Ranges of exclusion limits for gluinos and squarks, varying  $m(\tilde{\chi}^0)$




For limits on  $m(\tilde{g}), m(\tilde{q}) > m(\tilde{g})$  (and vice versa).  $\sigma^{\text{prod}} = \sigma^{\text{NLO-QCD}}$ .

$m(\tilde{\chi}^\pm), m(\tilde{\chi}_2^0) \equiv \frac{m(\tilde{g}) + m(\tilde{\chi}^0)}{2}$ .

$m(\tilde{\chi}^0)$  is varied from 0 GeV/c<sup>2</sup> (dark blue) to  $m(\tilde{g}) - 200$  GeV/c<sup>2</sup> (light blue).

# CMS Preliminary

Ranges of exclusion limits for gluinos and squarks, varying  $m(\tilde{\chi}^0)$



Limits on colored SUSY production are pretty much on track.

For limits on  $m(\tilde{g}), m(\tilde{q}) > m(\tilde{g})$  (and vice versa).  $\sigma^{\text{prod}} = \sigma^{\text{NLO-QCD}}$ .


$m(\tilde{\chi}^\pm), m(\tilde{\chi}_2^0) \equiv \frac{m(\tilde{g}) + m(\tilde{\chi}^0)}{2}$ .

$m(\tilde{\chi}^0)$  is varied from 0 GeV/c<sup>2</sup> (dark blue) to  $m(\tilde{g}) - 200$  GeV/c<sup>2</sup> (light blue).



# CMS Preliminary

Ranges of exclusion limits for gluinos and squarks, varying  $m(\tilde{\chi}^0)$



Limits on colored SUSY production are pretty much on track.

Squeezed spectra are currently one exception.


For limits on  $m(\tilde{g}), m(\tilde{q}) \gg m(\tilde{g})$  (and vice versa).  $\sigma^{\text{prod}} = \sigma^{\text{NLO-QCD}}$ .

$m(\tilde{\chi}^\pm), m(\tilde{\chi}_2^0) \equiv \frac{m(\tilde{g}) + m(\tilde{\chi}^0)}{2}$ .

$m(\tilde{\chi}^0)$  is varied from 0 GeV/c<sup>2</sup> (dark blue) to  $m(\tilde{g}) - 200$  GeV/c<sup>2</sup> (light blue).




# Squeezed Spectra



No limit on gluino mass  
for  $m_{\text{LSP}} > 400$  GeV ??


Keep in mind, squeezed spectra are perfectly valid theoretically!!  
They only do not arise in straw-man models like the CMSSM.

# ATLAS SUSY Searches\* - 95% CL Lower Limits (Status: March 2012)






# ATLAS SUSY Searches\* - 95% CL Lower Limits (Status: March 2012)




Limits on EW production currently weak,  
nearly non-existent.

# LHC Limits on EW production

Currently need to assume best-case scenarios to get a limit:



neutralino LSP with 100% BR to leptons




gravitino LSP with 100% BR to photons

# Identity of the LOSP: AMSB



- A general feature of anomaly mediation:
- $M_1:M_2:M_3 = 2.8 : 1 : 7$
- Wino LSP!!
- In AMSB, squarks and sleptons either  $\sim 100$  TeV (not sequestered), or weak scale (sequestered). Latter case leads to **tachyonic slepton masses** for pure AMSB.
- Version of “minimal AMSB” implemented in spectrum generators adds a universal scalar mass-squared to AMSB masses. **THIS IS NOT A REAL MODEL.** As problematic as the CMSSM.

# Identity of the LOSP: AMSB



- Charged wino can have a macroscopic lifetime before decaying to the neutral wino -- **disappearing track**
- ATLAS has a recent search (ATLAS-CONF-2012-034).
- Final state:  $\geq 3$  jets + lepton veto + MET + disappearing track
- It's a good start, but very non-inclusive!!! Highly tuned to specifics of minimal AMSB model.


# ATLAS search for disappearing tracks + (...)



Misleading plot!  
 Search is actually looking for gluino  
 and squark production...

# Identity of the LOSP: GMSB


- In GMSB, the LOSP (aka the NLSP) can be any superpartner in the MSSM. (Meade, Seiberg & DS; Buican, Meade, Seiberg, DS)





# Identity of the LOSP: GMSB

Analysis	Collaboration	Luminosity (fb <sup>-1</sup> )	Ref
jets+MET <i>with <math>\alpha_T</math></i>	ATLAS	1	[2]
	CMS	1.1	[3]
	CMS	1.1	[4]
6-8 jets+MET	ATLAS	1.34	[5]
<i>b</i> -jets+MET	ATLAS	0.833	[6]
	CMS	1.1	[7]
SS dileptons+jets+MET	CMS	0.98	[8]
OS dileptons +jets+ MET	CMS	0.98	[9]
lepton+jets+MET	ATLAS	1.04	[10]
	CMS	1.1	[11]
lepton+ <i>b</i> -jets+MET	ATLAS	1.03	[12]
$Z(\ell^+\ell^-)$ +jets+MET	CMS	0.98	[13]
$t\bar{t}$ +MET	ATLAS	1.04	[14]
$\gamma\gamma$ +MET	ATLAS	1.07	[15]
$\gamma\gamma$ +jet+MET	CMS	1.1	[16]
$\gamma$ +jets+MET	CMS	1.1	[16]
$\gamma+\ell$ +MET	CMS	0.035	[17]





(from Kats, Meade, Reece, DS; “Status of GMSB after 1/fb at the LHC”) Gluino mass [GeV]

- Overall, the coverage of different NLSP types by **non-dedicated** searches has been pretty good.
- Indicates that the search for SUSY at the LHC is robust. Well done!
- Again, limits are for strong production; limits on EW production still lacking.

# Identity of the LOSP: GMSB

- We're also starting to see more **dedicated searches** for general GMSB signatures.
- There's more to GMSB than photons and taus!!!
- For example, hot off the presses is an ATLAS search for **higgsino-like NLSPs decaying to Z+MET** ([ATLAS-CONF-2012-047](#))
- First ever dedicated search for higgsino NLSPs!
- $\text{gluino} \rightarrow \text{higgsino} + \text{jets}$ ;  $\text{higgsino} \rightarrow \text{Z} + \text{gravitino}$
- Final state:  $\text{Z}(\ell\ell) + \text{MET} + (\geq 3 \text{ jets or HT})$


# ATLAS Search for Higgsino NLSPs



Interestingly, no limit yet on direct Higgsino production...



# Higgsino NLSPs through Stop Production

- Another recent, related search from ATLAS: Higgsino NLSPs through stop production -- “natural SUSY” ([arXiv:1204.6736](https://arxiv.org/abs/1204.6736))
- $\text{Stop} \rightarrow b + \text{Higgsino}$  or  $t + \text{Higgsino}$ ;  $\text{Higgsino} \rightarrow Z + \text{MET}$
- Final state:  $Z(\ell) + \text{MET} + b \text{ jet} + \text{jets}$



# An Interesting Alternative: Stop NLSPs


- An even more minimal realization of “natural SUSY” is stop NLSP  
(Kats & DS “Light Stop NLSPs at the Tevatron and LHC”)
- Direct production of stop; stop  $\rightarrow$  top+MET



Very challenging to see under  $t\bar{t}$  background!

# An Interesting Alternative: Stop NLSPs

- Currently no dedicated searches for stop NLSPs, at either Tevatron or LHC. **They could still be lighter than the top!!**



(Kats & DS; Kats, Meade, Reece, DS)

# Conclusions

- “The state of the SUSY search at the LHC is strong.”
- Colored SUSY production has been well covered.
- Searches are mostly robust, not overly tuned to specific scenarios.
- The low hanging fruit has largely been picked.
- New, more challenging frontiers await:
  - EW production (winos, higgsinos, sleptons)
  - top-rich signatures (light stops)
  - Displaced vertices (intermediate SUSY-breaking scale)

**The End**